

A COUPLED FIXED POINT RESULT IN PARTIALLY ORDERED PARTIAL METRIC SPACES THROUGH IMPLICIT FUNCTION

Selma Gülyaz ^{*}, Erdal Karapınar [†]

Received 16:06:2012 : Accepted 09:11:2012

Abstract

In this manuscript, we discuss the existence of coupled fixed points in the context of partially ordered metric spaces through implicit relations for mappings $F : X \times X \rightarrow X$ such that F has the mixed monotone property. Our main theorem improves and extends various results in the literature. We also state an example to illustrate our work.

Keywords: Coupled fixed point; Mixed monotone property; Implicit relation; Ordered partial metric space

2000 AMS Classification: 47H10,54H25,46J10, 46J15

1. Introduction and Preliminaries

The notion of partial metric space, introduced by Matthews [46], is a generalization of metric space defined by Fréchet [20] in 1906. Roughly speaking the most remarkable property in a partial metric space is that the self-distance need not be zero. Nonzero self-distance makes perfect sense in the framework of Computer Sciences, in particular, in the Domain Theory and Semantics (see e.g., [39, 40, 51, 23, 57, 65, 66, 74, 75]). In the paper [46], Matthews proved an analog of the well-known Banach contraction principle in the context of complete partial metric spaces. After this result, many authors have conducted further research on fixed point theorems in the same class of spaces. Furthermore they studied topological properties of partial metric spaces (see e.g., [2]-[7], [9, 16, 18, 24, 25] [35]-[33],[40],[60]-[64],[67, 73]).

A partial metric is a function $p : X \times X \rightarrow [0, \infty)$ which satisfies the following conditions

(P1) $p(x, y) = p(y, x)$,

(P2) If $p(x, x) = p(x, y) = p(y, y)$, then $x = y$,

^{*}Department of Mathematics, Cumhuriyet University, Sivas, Turkey Email: sgulyaz@cumhuriyet.edu.tr

[†]Department of Mathematics, Atilim University 06836, İncek, Ankara, Turkey Email: ekarapinar@atilim.edu.tr

(P3) $p(x, x) \leq p(x, y)$,

(P4) $p(x, z) + p(y, y) \leq p(x, y) + p(y, z)$,

for all $x, y, z \in X$. Then the pair (X, p) is called a partial metric space.

1.1. Example. (See [68]) Let (X, d) and (X, p) be a metric space and a partial metric space, respectively. Mappings $\rho_i : X \times X \rightarrow \mathbb{R}^+$ for $i \in \{1, 2, 3\}$ defined by

$$\begin{aligned}\rho_1(x, y) &= d(x, y) + p(x, y) \\ \rho_2(x, y) &= d(x, y) + \max\{\omega(x), \omega(y)\} \\ \rho_3(x, y) &= d(x, y) + a\end{aligned}$$

induce partial metrics on X , where $\omega : X \rightarrow \mathbb{R}^+$ is an arbitrary function and $a \geq 0$.

Each partial metric p on X generates a T_0 topology τ_p on X with the family of open p -balls $\{B_p(x, \varepsilon) : x \in X, \varepsilon > 0\}$ as a base, where $B_p(x, \varepsilon) = \{y \in X : p(x, y) < p(x, x) + \varepsilon\}$ for all $x \in X$. Similarly, a closed p -ball is defined as $B_p[x, \varepsilon] = \{y \in X : p(x, y) \leq p(x, x) + \varepsilon\}$.

1.2. Definition. (See [46]) Let (X, p) be a partial metric space.

- (i) A sequence $\{x_n\}$ in X converges to $x \in X$ whenever $\lim_{n \rightarrow \infty} p(x, x_n) = p(x, x)$,
- (ii) A sequence $\{x_n\}$ in X is called Cauchy whenever $\lim_{n, m \rightarrow \infty} p(x_n, x_m)$ exists (and finite),
- (iii) (X, p) is said to be complete if every Cauchy sequence $\{x_n\}$ in X converges, with respect to τ_p , to a point $x \in X$, such that, $\lim_{n, m \rightarrow \infty} p(x_n, x_m) = p(x, x)$.
- (iv) A mapping $f : X \rightarrow X$ is said to be continuous at $x_0 \in X$ if for each $\varepsilon > 0$ there exists $\delta > 0$ such that $f(B(x_0, \delta)) \subset B(f(x_0), \varepsilon)$.

Romaguera (see [60]) defined the notion of 0-Cauchy sequence in partial metric spaces. Furthermore, he introduced the concept of 0-completeness in the same class of spaces.

1.3. Remark. Notice that the limit of a sequence in partial metric space need not to be unique. For more details and examples see e.g. [37].

1.4. Definition. (See [60]) A sequence $\{x_n\}$ in a partial metric space (X, p) is called 0-Cauchy if $\lim_{n, m \rightarrow \infty} p(x_n, x_m) = 0$. A partial metric space (X, p) is said to be 0-complete if every 0-Cauchy sequence in X converges, with respect to τ_p , to a point $x \in X$ such that $p(x, x) = 0$. In this case, p is said to be a 0-complete partial metric on X .

Observe that each 0-Cauchy sequence is also a Cauchy sequence in a partial metric space. In particular, we note that each complete partial metric is a 0-complete partial metric on X . However the converse is not true and the following example demonstrate that there exists a 0-complete partial metric that is not complete.

1.5. Example. (See [60, 61]) Let $(\mathbb{Q} \cap [0, \infty), p)$ be the partial metric space, where \mathbb{Q} and $p(x, y)$ represent the set of rational numbers and the partial metric $\max\{x, y\}$, respectively.

One of the characterizations of continuity of mappings in partial metric spaces was given by Samet et al. [64] as follows:

1.6. Lemma. (See [64]) Let (X, p) be a partial metric space. The function $F : X \rightarrow X$ is continuous if given a sequence $\{x_n\} \in \mathbb{N}$ and $x \in X$ such that $p(x, x) = \lim_{n \rightarrow +\infty} p(x, x_n)$, then

$$p(Fx, Fx) = \lim_{n \rightarrow +\infty} p(Fx, Fx_n).$$

1.7. Example. (See [64]) Consider $X = [0, \infty)$ endowed with the partial metric $p : X \times X \rightarrow [0, \infty)$ defined by $p(x, y) = \max\{x, y\}$ for all $x, y \geq 0$. Let $F : X \rightarrow X$ be a non-decreasing function. If F is continuous with respect to the standard metric $d(x, y) = |x - y|$ for all $x, y \geq 0$, then F is continuous with respect to the partial metric p .

There is a close relationship between metrics and partial metrics. Indeed, if p is a partial metric on X , then the function $d_p : X \times X \rightarrow [0, \infty)$ given by

$$d_p(x, y) = 2p(x, y) - p(x, x) - p(y, y)$$

is a metric on X . Moreover,

$$(1.1) \quad \lim_{n \rightarrow \infty} d_p(x, x_n) = 0 \Leftrightarrow \lim_{n \rightarrow \infty} p(x, x_n) = \lim_{n, m \rightarrow \infty} p(x_n, x_m) = p(x, x).$$

1.8. Lemma. (See e.g. [46]) Let (X, p) be a partial metric space.

- (a) A sequence $\{x_n\}$ is Cauchy if and only if $\{x_n\}$ is a Cauchy sequence in the metric space (X, d_p) ,
- (b) (X, p) is complete if and only if the metric space (X, d_p) is complete.

The following lemmas will be frequently used in the proof of the main result.

1.9. Lemma. Let (X, p) be a partial metric space. Then

- (a) If $p(x, y) = 0$, then $x = y$,
- (b) If $x \neq y$, then $p(x, y) > 0$.
- (c) If $x_n \rightarrow z$ with $p(z, z) = 0$, then $\lim_{n \rightarrow \infty} p(x_n, y) = p(z, y)$ for all $y \in X$.

In [26], Guo and Lakshmikantham introduced the notion of coupled fixed point. In 2006, Gnana-Bhaskar and Lakshmikantham[10] defined the notion of mixed monotone mapping and reconsidered coupled fixed point in the context of partially ordered set. In this initial paper, the authors proved some coupled fixed point theorems for the mixed monotone mappings and discussed the existence and uniqueness of solution for a periodic boundary value problem. Next, we recall the necessary definitions and their basic results in this direction.

1.10. Definition. (See [10]) Let (X, \preceq) be a partially ordered set. The mapping $F : X \times X \rightarrow X$ is said to have the mixed monotone property if $F(x, y)$ is monotone non-decreasing in x and is monotone non-increasing in y , that is, for any $x, y \in X$,

$$x_1, x_2 \in X, \quad x_1 \preceq x_2 \Rightarrow F(x_1, y) \preceq F(x_2, y)$$

and

$$y_1, y_2 \in X, \quad y_1 \preceq y_2 \Rightarrow F(x, y_1) \succeq F(x, y_2).$$

1.11. Definition. (See [10]) An element $(x, y) \in X \times X$ is called a coupled fixed point of the mapping $F : X \times X \rightarrow X$ if

$$x = F(x, y) \text{ and } y = F(y, x).$$

We state now the main results of Gnana-Bhaskar and Lakshmikantham in [10].

1.12. Theorem. (See [10]) Let (X, \preceq) be a partially ordered set. Suppose there exists a metric d on X such that (X, d) is a complete metric space. Assume that there exists a $k \in [0, 1)$ with

$$d(F(x, y), F(u, v)) \leq \frac{k}{2} [d(x, u) + d(y, v)]$$

for all $x \succeq u$ and $y \preceq v$. Let either

- (a) $F : X \times X \rightarrow X$ be a continuous mapping with the mixed monotone property on X , or,

(b) X has the following property:

- (i) if a non-decreasing sequence $\{x_n\} \rightarrow x$, then $x_n \preceq x$ for all n ,
- (ii) if a non-increasing sequence $\{y_n\} \rightarrow y$, then $y \preceq y_n$ for all n .

If there exist two elements $x_0, y_0 \in X$ with

$$x_0 \preceq F(x_0, y_0) \quad \text{and} \quad y_0 \succeq F(y_0, x_0)$$

then there exist $x, y \in X$ such that

$$x = F(x, y) \quad \text{and} \quad y = F(y, x).$$

Following Theorem 1.12, several coupled coincidence/fixed point theorems and their applications to integral equations, matrix equations, periodic boundary value problems have been reported (see e.g. [9, 11, 12, 13, 21, 38, 29, 34, 41, 43, 44, 42, 45, 63] and references therein).

In nonlinear analysis, especially in fixed point theory, implicit relations on metric spaces have been investigated heavily in many articles (see, e.g., [6], [19, 22], [53] and references therein). In this paper, by using implicit relations, we examine the existence of a coupled fixed point theorem for mappings $F : X \times X \rightarrow X$ satisfying the mixed monotone property in the context of partial metric spaces. A set of implicit relations, denoted by \mathbb{H} , is the collection of all continuous functions $H : (\mathbb{R}^+)^5 \rightarrow \mathbb{R}$ which satisfy

- (H1) $H(t_1, t_2, t_3, t_4, t_5)$ is non-increasing in t_2 and t_5 , and
- (H2) there exists a function $\varphi \in \Phi$ such that

$$H(u, u + v, v, w, u + v) \leq 0 \quad \text{implies} \quad u \leq \varphi(\max\{v, w\})$$

where Φ denotes the set of all functions $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with the properties

- (i) φ is continuous and non-decreasing,
- (ii) $\varphi(t) < t$ for each $t > 0$ and $\varphi(0) = 0$.

1.13. Example. It is easy to check that the following functions are in \mathbb{H}

- (H₁) $H(t_1, t_2, t_3, t_4, t_5) = t_1 - \alpha t_2 - \beta t_3 - \gamma t_4 - \theta t_5$, where $\alpha, \beta, \gamma, \theta$ are non-negative real numbers satisfying $2\alpha + \beta + \gamma + 2\theta < 1$.
- (H₂) $H(t_1, t_2, t_3, t_4, t_5) = t_1 - \alpha \max\{t_2/2, t_3, t_4, t_5/2\}$, where $\alpha \in (0, 1)$.
- (H₃) $H(t_1, t_2, t_3, t_4, t_5) = t_1 - \varphi(\max\{t_3, t_4\})$, where $\varphi \in \Phi$.

In this paper, we prove a coupled fixed point theorem for mappings satisfying certain implicit relations in the framework of partial metric spaces.

2. Coupled fixed point theorem

We start this section with our main result.

2.1. Theorem. Let (X, p, \preceq) be a partially ordered complete partial metric space. Suppose $F : X \times X \rightarrow X$ be a mapping such that F has the mixed monotone property. Assume that there exists $H \in \mathbb{H}$ such that

$$(2.1) \quad H \left(\begin{array}{c} p(F(x, y), F(u, v)), p(F(x, y), x) + p(F(u, v), u), \\ p(x, u), p(y, v), p(F(x, y), u) + p(F(u, v), x) \end{array} \right) \leq 0$$

for all $x, y, u, v \in X$ with $x \succeq u$ and $y \preceq v$. Suppose that either

- (a) F is continuous or
- (b) X has the following property
 - (i) if a non-decreasing sequence $\{x_n\} \rightarrow x$, then $x_n \preceq x$ for all n ,
 - (ii) if a non-increasing sequence $\{y_n\} \rightarrow y$, then $y \preceq y_n$ for all n .

If there exist two elements $x_0, y_0 \in X$ with

$$x_0 \preceq F(x_0, y_0) \text{ and } y_0 \succeq F(y_0, x_0),$$

then F has a coupled fixed point in X .

Proof. Let $x_0, y_0 \in X$ be such that $x_0 \preceq F(x_0, y_0)$ and $y_0 \succeq F(y_0, x_0)$. We construct the iterative sequences $\{x_n\}$ and $\{y_n\}$ in X as follows

$$(2.2) \quad x_{n+1} = F(x_n, y_n) \text{ and } y_{n+1} = F(y_n, x_n) \text{ for all } n \geq 0.$$

By using the mathematical induction and the mixed monotone property of F , we can show that

$$(2.3) \quad x_n \preceq x_{n+1} \text{ and } y_n \succeq y_{n+1} \text{ for all } n \geq 0.$$

If there is some $n_0 \in \mathbb{N}^*$ such that $x_{n_0} = x_{n_0+1}$ and $y_{n_0} = y_{n_0+1}$ then

$$x_{n_0} = x_{n_0+1} = F(x_{n_0}, y_{n_0}) \text{ and } y_{n_0} = y_{n_0+1} = F(y_{n_0}, x_{n_0})$$

which concludes that (x_{n_0}, y_{n_0}) is a coupled fixed point of F . Thus we assume that $x_{n_0} \neq x_{n_0+1}$ or $y_{n_0} \neq y_{n_0+1}$ for all n . By Lemma 1.9, we have $\max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\} > 0$ for all n .

Since $x_{n+1} \succeq x_n$ and $y_{n+1} \preceq y_n$, from (2.1), we have

$$H \left(\begin{array}{c} p(F(x_{n+1}, y_{n+1}), F(x_n, y_n)), p(F(x_{n+1}, y_{n+1}), x_{n+1}) + p(F(x_n, y_n), x_n), \\ p(x_{n+1}, x_n), p(y_{n+1}, y_n), p(F(x_{n+1}, y_{n+1}), x_n) + p(F(x_n, y_n), x_{n+1}) \end{array} \right) \leq 0$$

or

$$(2.4) \quad H \left(\begin{array}{c} p(x_{n+2}, x_{n+1}), p(x_{n+2}, x_{n+1}) + p(x_{n+1}, x_n), \\ p(x_{n+1}, x_n), p(y_{n+1}, y_n), p(x_{n+2}, x_n) + p(x_{n+1}, x_{n+1}) \end{array} \right) \leq 0.$$

Due to (P4) we have,

$$(2.5) \quad \begin{aligned} p(x_{n+2}, x_n) &\leq p(x_{n+2}, x_{n+1}) + p(x_{n+1}, x_n) - p(x_{n+1}, x_{n+1}) \\ &\leq p(x_{n+2}, x_{n+1}) + p(x_{n+1}, x_n). \end{aligned}$$

By the properties of H and (2.5), the inequality in (2.4) turns into

$$H \left(\begin{array}{c} p(x_{n+2}, x_{n+1}), p(x_{n+2}, x_{n+1}) + p(x_{n+1}, x_n), \\ p(x_{n+1}, x_n), p(y_{n+1}, y_n), p(x_{n+2}, x_{n+1}) + p(x_{n+1}, x_n) \end{array} \right) \leq 0$$

which yields that

$$(2.6) \quad p(x_{n+2}, x_{n+1}) \leq \varphi(\max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\}).$$

Similarly, one can show that

$$(2.7) \quad p(y_{n+2}, y_{n+1}) \leq \varphi(\max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\}).$$

From (2.6) and (2.7), we have

$$(2.8) \quad \max\{p(x_{n+2}, x_{n+1}), p(y_{n+2}, y_{n+1})\} \leq \varphi(\max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\})$$

which implies

$$\max\{p(x_{n+2}, x_{n+1}), p(y_{n+2}, y_{n+1})\} < \max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\}$$

by the property of φ . This means that $\{p_n := \max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\}\}$ is a decreasing sequence of positive real numbers. So there is an $L \geq 0$ such that

$$(2.9) \quad \lim_{n \rightarrow \infty} p_n = \lim_{n \rightarrow \infty} \max\{p(x_{n+1}, x_n), p(y_{n+1}, y_n)\} = L.$$

We shall show that $L = 0$. Assume, to the contrary, that $L > 0$. Taking $n \rightarrow \infty$ in (2.8), we have

$$L \leq \lim_{n \rightarrow \infty} \varphi(p_n) = \varphi(L) < L$$

which is a contradiction. Thus $L = 0$. Hence,

$$(2.10) \quad \lim_{n \rightarrow \infty} p(x_{n+1}, x_n) = 0 \text{ and } \lim_{n \rightarrow \infty} p(y_{n+1}, y_n) = 0.$$

Next, we show that $\{x_n\}$ and $\{y_n\}$ are Cauchy sequences. Suppose, to the contrary, that at least one of $\{x_n\}$ or $\{y_n\}$ is not a Cauchy sequence. This means that there exists an $\varepsilon > 0$ for which we can find subsequences $\{x_{n(k)}\}$, $\{x_{m(k)}\}$ of $\{x_n\}$ and $\{y_{n(k)}\}$, $\{y_{m(k)}\}$ of $\{y_n\}$ with $n(k) > m(k) \geq k$ such that

$$(2.11) \quad \max \{p(x_{n(k)}, x_{m(k)}), p(y_{n(k)}, y_{m(k)})\} \geq \varepsilon.$$

Furthermore, corresponding to $m(k)$, we can choose $n(k)$ in such a way that it is the smallest integer with $n(k) > m(k) \geq k$ and satisfies (2.11). Then

$$(2.12) \quad \max \{p(x_{n(k)-1}, x_{m(k)}), p(y_{n(k)-1}, y_{m(k)})\} < \varepsilon.$$

Using the triangle inequality and (2.12), we have

$$(2.13) \quad \begin{aligned} p(x_{n(k)}, x_{m(k)}) &\leq p(x_{n(k)}, x_{n(k)-1}) + p(x_{n(k)-1}, x_{m(k)}) - p(x_{n(k)-1}, x_{n(k)-1}) \\ &\leq p(x_{n(k)}, x_{n(k)-1}) + p(x_{n(k)-1}, x_{m(k)}) \\ &< p(x_{n(k)}, x_{n(k)-1}) + \varepsilon \end{aligned}$$

and

$$(2.14) \quad \begin{aligned} p(y_{n(k)}, y_{m(k)}) &\leq p(y_{n(k)}, y_{n(k)-1}) + p(y_{n(k)-1}, y_{m(k)}) - p(y_{n(k)-1}, y_{n(k)-1}) \\ &\leq p(y_{n(k)}, y_{n(k)-1}) + p(y_{n(k)-1}, y_{m(k)}) \\ &< p(y_{n(k)}, y_{n(k)-1}) + \varepsilon. \end{aligned}$$

From (2.11), (2.13) and (2.14), we have

$$\begin{aligned} \varepsilon &\leq \max \{p(x_{n(k)}, x_{m(k)}), p(y_{n(k)}, y_{m(k)})\} \\ &< \max \{p(x_{n(k)}, x_{n(k)-1}), p(y_{n(k)}, y_{n(k)-1})\} + \varepsilon. \end{aligned}$$

Letting $k \rightarrow \infty$ in the inequality above and using (2.9) we get

$$(2.15) \quad \lim_{k \rightarrow \infty} \max \{p(x_{n(k)}, x_{m(k)}), p(y_{n(k)}, y_{m(k)})\} = \varepsilon.$$

By the triangle inequality

$$\begin{aligned} p(x_{n(k)}, x_{m(k)}) &\leq p(x_{n(k)}, x_{n(k)-1}) + p(x_{n(k)-1}, x_{m(k)-1}) + p(x_{m(k)-1}, x_{m(k)}) \\ &\quad - p(x_{n(k)-1}, x_{n(k)-1}) - p(x_{m(k)-1}, x_{m(k)-1}) \\ &\leq p(x_{n(k)}, x_{n(k)-1}) + p(x_{n(k)-1}, x_{m(k)-1}) + p(x_{m(k)-1}, x_{m(k)}) \end{aligned}$$

and

$$\begin{aligned} p(y_{n(k)}, y_{m(k)}) &\leq p(y_{n(k)}, y_{n(k)-1}) + p(y_{n(k)-1}, y_{m(k)-1}) + p(y_{m(k)-1}, y_{m(k)}) \\ &\quad - p(y_{n(k)-1}, y_{n(k)-1}) - p(y_{m(k)-1}, y_{m(k)-1}) \\ &\leq p(y_{n(k)}, y_{n(k)-1}) + p(y_{n(k)-1}, y_{m(k)-1}) + p(y_{m(k)-1}, y_{m(k)}). \end{aligned}$$

From the last two inequalities and (2.11), we have

$$(2.16) \quad \begin{aligned} \varepsilon &\leq \max \{p(x_{n(k)}, x_{m(k)}), p(y_{n(k)}, y_{m(k)})\} \\ &\leq \max \{p(x_{n(k)}, x_{n(k)-1}), p(y_{n(k)}, y_{n(k)-1})\} \\ &\quad + \max \{p(x_{m(k)-1}, x_{m(k)}), p(y_{m(k)-1}, y_{m(k)})\} \\ &\quad + \max \{p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1})\}. \end{aligned}$$

Again, by the triangle inequality,

$$\begin{aligned} p(x_{n(k)-1}, x_{m(k)-1}) &\leq p(x_{n(k)-1}, gx_{m(k)}) + p(x_{m(k)}, x_{m(k)-1}) - p(x_{m(k)}, x_{m(k)}) \\ &\leq p(x_{n(k)-1}, x_{m(k)}) + p(x_{m(k)}, x_{m(k)-1}) \\ &< p(x_{m(k)}, x_{m(k)-1}) + \varepsilon \end{aligned}$$

and

$$\begin{aligned} p(y_{n(k)-1}, y_{m(k)-1}) &\leq p(y_{n(k)-1}, gy_{m(k)}) + p(y_{m(k)}, y_{m(k)-1}) - p(y_{m(k)}, y_{m(k)}) \\ &\leq p(y_{n(k)-1}, y_{m(k)}) + p(y_{m(k)}, y_{m(k)-1}) \\ &< p(y_{m(k)}, y_{m(k)-1}) + \varepsilon. \end{aligned}$$

Therefore,

$$(2.17) \quad \max\{p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1})\} < \max\{p(x_{m(k)}, x_{m(k)-1}), p(y_{m(k)}, y_{m(k)-1})\} + \varepsilon.$$

From (2.16) and (2.17), we have

$$\begin{aligned} \varepsilon &- \max\{p(x_{n(k)}, x_{n(k)-1}), p(y_{n(k)}, y_{n(k)-1})\} \\ &- \max\{p(x_{m(k)-1}, x_{m(k)}), p(y_{m(k)-1}, y_{m(k)})\} \\ &\leq \max\{p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1})\} \\ &< \max\{p(x_{m(k)}, x_{m(k)-1}), p(y_{m(k)}, y_{m(k)-1})\} + \varepsilon. \end{aligned}$$

Taking the limit as $k \rightarrow \infty$ in the inequalities above and using (2.9), we get

$$(2.18) \quad \lim_{k \rightarrow \infty} \max\{p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1})\} = \varepsilon.$$

From (2.15) and (2.18), the sequences $\{p(x_{n(k)}, x_{m(k)})\}$, $\{p(y_{n(k)}, y_{m(k)})\}$, $\{p(x_{n(k)-1}, x_{m(k)-1})\}$ and $\{p(y_{n(k)-1}, y_{m(k)-1})\}$ have subsequences converging to $\varepsilon_1, \varepsilon_2, \varepsilon_3$ and ε_4 , respectively, and $\max\{\varepsilon_1, \varepsilon_2\} = \max\{\varepsilon_3, \varepsilon_4\} = \varepsilon > 0$. We may assume that

$$\begin{aligned} \lim_{k \rightarrow \infty} p(x_{n(k)}, x_{m(k)}) &= \varepsilon_1, & \lim_{k \rightarrow \infty} p(y_{n(k)}, y_{m(k)}) &= \varepsilon_2, \\ \lim_{k \rightarrow \infty} p(x_{n(k)-1}, x_{m(k)-1}) &= \varepsilon_3 \text{ and } \lim_{k \rightarrow \infty} p(y_{n(k)-1}, y_{m(k)-1}) &= \varepsilon_4. \end{aligned}$$

We first suppose that $\varepsilon_1 = \max\{\varepsilon_1, \varepsilon_2\} = \varepsilon$. Since $n(k) > m(k)$, $x_{n(k)-1} \succeq x_{m(k)-1}$ and $y_{n(k)-1} \preceq y_{m(k)-1}$. From (2.1), we have

$$H \left(\begin{array}{l} p(F(x_{n(k)-1}, y_{n(k)-1}), F(x_{m(k)-1}, y_{m(k)-1})), p(F(x_{n(k)-1}, y_{n(k)-1}), x_{n(k)-1}) + \\ p(F(x_{m(k)-1}, y_{m(k)-1}), x_{m(k)-1}), p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1}), \\ p(F(x_{n(k)-1}, y_{n(k)-1}), x_{m(k)-1}) + p(F(x_{m(k)-1}, y_{m(k)-1}), x_{n(k)-1}) \end{array} \right) \leq 0$$

or

$$H \left(\begin{array}{l} p(x_{n(k)}, x_{m(k)}), p(x_{n(k)}, x_{n(k)-1}) + p(x_{m(k)}, x_{m(k)-1}), \\ p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1}), p(x_{n(k)}, x_{m(k)-1}) + p(x_{m(k)}, x_{n(k)-1}) \end{array} \right) \leq 0$$

or

$$H \left(\begin{array}{l} p(x_{n(k)}, x_{m(k)}), p(x_{n(k)}, x_{n(k)-1}) + p(x_{m(k)}, x_{m(k)-1}), \\ p(x_{n(k)-1}, x_{m(k)-1}), p(y_{n(k)-1}, y_{m(k)-1}), p(x_{n(k)}, x_{m(k)}) + \\ p(x_{m(k)}, x_{m(k)-1}) + p(x_{m(k)}, x_{m(k)-1}) + p(x_{m(k)-1}, x_{n(k)-1}) \end{array} \right) \leq 0.$$

Letting $k \rightarrow \infty$ in the last inequality together with (2.10), we derive

$$H(\varepsilon_1, 0, \varepsilon_3, \varepsilon_4, \varepsilon_1 + \varepsilon_3) \leq 0.$$

Hence, we get

$$H(\varepsilon_1, \varepsilon_1 + \varepsilon_3, \varepsilon_3, \varepsilon_4, \varepsilon_1 + \varepsilon_3) \leq 0$$

which implies $\varepsilon = \varepsilon_1 \leq \varphi(\max\{\varepsilon_3, \varepsilon_4\}) = \varphi(\varepsilon) < \varepsilon$. This is a contradiction.

Using the same argument as above for the case $\varepsilon_2 = \max\{\varepsilon_1, \varepsilon_2\} = \varepsilon$, we also get a contradiction. Thus $\{x_n\}$ and $\{y_n\}$ are Cauchy sequences. Since X is complete, there exist $x, y \in X$ such that

$$(2.19) \quad \begin{aligned} \lim_{n,m \rightarrow \infty} p(x_n, x_m) &= \lim_{n \rightarrow \infty} p(x_n, x) = p(x, x) \\ &\text{and} \\ \lim_{n,m \rightarrow \infty} p(y_n, y_m) &= \lim_{n \rightarrow \infty} p(y_n, y) = p(y, y). \end{aligned}$$

We want to show that $p(x, x) = 0 = p(y, y)$. Suppose, on the contrary, that

$$(2.20) \quad p(x, x) = \delta > 0 \text{ and } p(y, y) = \gamma > 0.$$

Then we see that

$$H \left(\begin{array}{c} p(F(x_{n-1}, y_{n-1}), F(x_{m-1}, y_{m-1})), p(F(x_{n-1}, y_{n-1}), x_{n-1}) + \\ p(F(x_{m-1}, y_{m-1}), x_{m-1}), p(x_{n-1}, x_{m-1}), p(y_{n-1}, y_{m-1}), \\ p(F(x_{n-1}, y_{n-1}), x_{m-1}) + p(F(x_{m-1}, y_{m-1}), x_{n-1}) \end{array} \right) \leq 0$$

or

$$H \left(\begin{array}{c} p(x_n, x_m), p(x_n, x_{n-1}) + p(x_m, x_{m-1}), \\ p(x_{n-1}, x_{m-1}), p(y_{n-1}, y_{m-1}), p(x_n, x_{m-1}) + p(x_m, x_{n-1}) \end{array} \right) \leq 0.$$

By using the triangle inequality (P4), we get

$$H \left(\begin{array}{c} p(x_n, x_m), p(x_n, x_{n-1}) + p(x_m, x_{m-1}), \\ p(x_{n-1}, x_{m-1}), p(y_{n-1}, y_{m-1}), p(x_n, x_m) + \\ p(x_m, x_{m-1}) + p(x_m, x_{m-1}) + p(x_{m-1}, x_{n-1}) \end{array} \right) \leq 0.$$

Letting $k \rightarrow \infty$, we derive

$$H(\delta, 0, \delta, 0, \delta) \leq 0$$

by (2.10) and (2.20). Hence, we find

$$H(\delta, 0 + \delta, \delta, 0, \delta + 0) \leq H(\delta, 0, \delta, 0, \delta) \leq 0$$

which implies $\delta \leq \varphi(\max\{\delta, 0\}) = \varphi(\delta) < \delta$. This is a contradiction. Hence $\delta = 0$. Analogously we find that $\gamma = 0$.

Now, suppose that the assumption (a) holds. We have

$$(2.21) \quad p(x, F((x_n), (y_n))) \leq p(x, F(x_n, y_n)) + p(F(x_n, y_n), F(x_n, y_n)).$$

Taking the limit as $n \rightarrow \infty$ in (2.21) and by (2.19), and the continuity of F we get $p(x, F(x, y)) = 0$.

Similarly, we can show that $p(y, F(y, x)) = 0$. Therefore, $x = F(x, y)$ and $y = F(y, x)$.

Finally, suppose that the assumption (b) holds. Since $\{x_n\}$ is a non-decreasing sequence and $x_n \rightarrow x$ and $\{y_n\}$ is a non-increasing sequence and $y_n \rightarrow y$, by the assumption, we have $x_n \preceq x$ and $y_n \succeq y$ for all n . Regarding (2.2) and (2.19), we have

$$(2.22) \quad \lim_{n \rightarrow \infty} p(x_n, x) = p(x, x) = \lim_{n \rightarrow \infty} p(F(x_n, y_n), x)$$

and

$$(2.23) \quad \lim_{n \rightarrow \infty} p(y_n, y) = p(y, y) = \lim_{n \rightarrow \infty} p(F(y_n, x_n), y).$$

We also have

$$H \left(\begin{array}{c} p(F(x_n, y_n), F(x, y)), p(F(x_n, y_n)) + p(F(x, y), x) \\ p(x_n, x), p(y_n, y), p(F(x_n, y_n), x) + p(F(x, y), x_n) \end{array} \right) \leq 0.$$

Letting $n \rightarrow \infty$ and using (2.22) and (2.23), we have

$$H(p(x, F(x, y)), p(x, F(x, y)), 0, 0, p(x, F(x, y))) \leq 0$$

which implies that $p(x, F(x, y)) \leq \varphi(\max\{0, 0\}) = 0$. Hence $x = F(x, y)$. Similarly, one can show that $y = F(y, x)$.

Thus we proved that F has a coupled fixed point in X . \square

2.2. Example. (See, e.g., [45]) Let $X = [0, \infty)$ with usual order \leq . Then, (X, p, \leq) be a partially ordered partial metric space where $p(x, y) = \max\{x, y\}$. Suppose $F(x, y) = \begin{cases} \frac{x-y}{2} & \text{if } x \geq y, \\ 0 & \text{otherwise.} \end{cases}$ and $H(t_1, t_2, t_3, t_4, t_5) = t_1 - \frac{1}{2} \max\{t_3, t_4\}$. It is clear that all conditions of Theorem 2.1 are satisfied. Notice that $(0, 0)$ is the coupled fixed point of the operator F .

References

- [1] Abbas, M., Sintunavarat, W. and Kumam, P. *Coupled fixed point in partially ordered G-metric spaces*, Fixed Point Theory and Applications **2012**, 2012:31.
- [2] Abbas, M., Nazir, T. and Romaguera, S. *Fixed point results for generalized cyclic contraction mappings in partial metric spaces*, Revista de la Real Academia de Ciencias Exactas, (in press), doi:10.1007/s13398-011-0051-5.
- [3] Abdeljawad, T. *Fixed Points for generalized weakly contractive mappings in partial metric spaces*, Math. Comput. Modelling **54**(11-12), 2923–2927, 2011.
- [4] Abdeljawad, T., Karapinar, E. and Taş, K. *Existence and uniqueness of common fixed point on partial metric spaces*, Appl. Math. Lett. **24**, 1894–1899, 2011.
- [5] Abdeljawad, T., Karapinar, E. and Taş, K. *A generalized contraction principle with control functions on partial metric spaces*, Comput. Math. Appl. **63** (3), 716–719, 2012.
- [6] Altun, I. and Turkoglu, D. *Some fixed point theorems for weakly compatible multivalued mappings satisfying an implicit relation*, Filomat **22**, 13–23, 2008.
- [7] Altun, I., Sola, F. and Simsek, H. *Generalized contractions on partial metric spaces*, Topology and Appl. **157** (18), 2778–2785, 2010.
- [8] Aydi, H., Vetro, C., Sintunavarat, W. and Kumam, P. *Coincidence and fixed points for contractions and cyclical contractions in partial metric spaces*, Fixed Point Theory and Applications, (to appear).
- [9] Aydi, H., Karapinar, E. and Shatanawi, W. *Coupled fixed point results for (ψ, φ) -weakly contractive condition in ordered partial metric spaces*, Comput. Math. Appl. **62**(12), 4449–4460, 2011.
- [10] Gnana Bhaskar, T. and Lakshmikantham, V. *Fixed point theorems in partially ordered metric spaces and applications*, Nonlinear Anal. **65**, 1379–1393, 2006.
- [11] Berinde, V. *Generalized coupled fixed point theorems for mixed monotone mappings in partially ordered metric spaces*, Nonlinear Anal. **74**, 7347–7355, 2011.
- [12] Berinde, V. *Coupled fixed point theorems for ϕ -contractive mixed monotone mappings in partially ordered metric spaces*, Nonlinear Anal. **75**, 3218–3228, 2012.
- [13] Choudhury, B.S. and Kundu, A. *A coupled coincidence point result in partially ordered metric spaces for compatible mappings*, Nonlinear Analysis **73**, 2524–2531, 2010.
- [14] Choudhury, B.S., Metiya, N. and Kundu, A. *Coupled coincidence point theorems in ordered metric spaces*. Ann. Univ. Ferrara **57**, 1–16, 2011.
- [15] Ćirić, L., Ćakic, N., Rajović, M. and Ume, J.S. *Monotone generalized nonlinear contractions in partially ordered metric spaces*, Fixed Point Theory Appl. **2008**, Art. ID 131294, 2008.
- [16] Chi, K.P., Karapinar, E. and Thanh, T. D. *A generalized contraction principle in partial metric spaces*, Math. Comput. Modelling **55** (5–6), 1673–1681, 2012.
- [17] Ćirić, Lj. *On contraction type mappings*, Math. Balkanica **1**, 52–57, 1971.
- [18] Ćirić, Lj., Samet, B., Aydi, H. and Vetro, C. *Common fixed points of generalized contractions on partial metric spaces and an application*, Appl. Math. Comput. **18** (6), 2398–2406, 2011.

- [19] Djoudi, A. and Aliouche, A. *A general common fixed point theorem for reciprocally continuous mappings satisfying an implicit relation*, The Austral. J. Math. Anal. Appl. **3**, 1–7, 2006.
- [20] Fréchet, M. *Sur quelques points du calcul fonctionnel*, Rend. Circ. Mat. Palermo **22**, 1–74, 1906.
- [21] Harjani, J., Lopez, B. and Sadarangani, K. *Fixed point theorems for mixed monotone operators and applications to integral equations*, Nonlinear Anal. **74**, 1749–1760, 2011.
- [22] Hung, N.M., Karapınar, E. and Luong, N.V. *Coupled coincidence point theorem in partially ordered metric spaces via implicit relation*, Abstract and Applied Analysis **2012**, Art. ID 796964, 2012.
- [23] Heckmann, R. *Approximation of metric spaces by partial metric spaces*, Applied Categorical Structures **7**, 71–83, 1999.
- [24] Ilić, D., Pavlović, V. and Rakočević, V. *Some new extensions of Banach's contraction principle to partial metric space*, Appl. Math. Lett. **24**(8), 1326–1330, 2011.
- [25] Ilić, D., Pavlović, V. and Rakočević, V. *Extensions of the Zamfirescu theorem to partial metric spaces* *Original Research Article*, Math. Comput. Modelling **55**(3–4), 801–809, 2012.
- [26] D.Guo, V. Lakshmikantham, *Coupled fixed points of nonlinear operators with applications*. Nonlinear Anal., 11 (1987) 623–632.
- [27] Jachymski, J. *Equivalent Conditions and the Meir-Keeler Type Theorems*, J.Math. Anal. Appl. **194** (1), 293–303, 1995.
- [28] Kadelburg, Z. and Radenović, S. *Meir-Keeler-type conditions in abstract metric spaces*, Appl.Math. Lett. **24** (8), 1411–1414, 2011.
- [29] Karapınar, E. *Coupled fixed point theorems for nonlinear contractions in cone metric spaces*, Computers and Mathematics with Applications **59**, 3656–3668, 2010.
- [30] Karapınar, E. and Erhan, I. M. *Fixed Point Theorems for Operators on Partial Metric Spaces*, Appl. Math. Lett. **24**, 1900–1904, 2011.
- [31] Karapınar, E. *Generalizations of Caristi Kirk's Theorem on Partial Metric Spaces*, Fixed Point Theory Appl. **2011**(4), doi:10.1186/1687-1812-2011-4, 2011.
- [32] Karapınar, E. and Yuksel, U. *Some common fixed point theorems in partial metric spaces*, Journal of Applied Mathematics **2011**, Art. ID 263621, 2011.
- [33] Karapınar, E. *A note on common fixed point theorems in partial metric spaces*, Miskolc Mathematical Notes **12** (2), 185–191, 2011.
- [34] Karapınar, E. *Couple Fixed Point on Cone Metric Spaces*, Gazi University Journal of Science **24**, 51–58, 2011.
- [35] Karapınar, E. *Weak ϕ -contraction on partial metric spaces*, J. Comput. Anal. Appl. **14** (2), 206–210, 2012.
- [36] Karapınar, E., Erhan, I.M. and Ulus, A.Y. *Fixed Point Theorem for Cyclic Maps on Partial Metric Spaces*, Appl. Math. Inf. Sci. **6** (1), 239–244, 2012.
- [37] Karapınar, E., Shobkolaei, N., Sedghi, S. and Vaezpour, S.M. *A common fixed point theorem for cyclic operators on partial metric spaces*, FILOMAT **26**(2), 407–414, 2012.
- [38] Karapınar, E., Nguyen Van Luong, Nguyen Xuan Thuan, Trinh Thi Hai, *Coupled coincidence points for mixed monotone operators in partially ordered metric spaces*, Arabian Journal of Mathematics, 1(2012),no: 3, 329–339.
- [39] Kopperman, R. D., Matthews, S. G. and Pajoohesh, H. *What do partial metrics represent?*, (Notes distributed at the 19th Summer Conference on Topology and its Applications, University of CapeTown, 2004).
- [40] Künzi, H. P. A., Pajoohesh, H. and Schellekens, M.P. *Partial quasi-metrics*, Theoretical Computer Science **365**(3), 237–246, 2006.
- [41] Lakshmikantham, V. and Ćirić, L. *Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces*, Nonlinear Anal. **70**, 4341–4349, 2009.
- [42] Luong, N. V. and Thuan, N. X. *Coupled fixed point theorems in partially ordered metric spaces*, Bull. Math. Anal. Appl. **2** (4), 16–24, 2010.
- [43] Luong, N. V. and Thuan, N. X. *Coupled fixed points in partially ordered metric spaces and application*, Nonlinear Anal. **74**, 983–992, 2011.
- [44] Luong, N. V. and Thuan, N. X. *Coupled fixed point theorems for mixed monotone mappings and an application to integral equations*, Compt. Math. Appl. **62**, 4238–4248, 2011.

- [45] Luong, N. V. and Thuan, N. X. *Coupled points in ordered generalized metric spaces and application to integro-differential equations*, (Submitted).
- [46] Matthews, S. G. *Partial metric topology*, in: *Proceedings 8th Summer Conference on General Topology and Applications*, Ann. New York Acad. Sci. **728**, 183–197, 1994.
- [47] Meir, A. and Keeler, E. *A theorem on contraction mapping*, J. Math. Anal. Appl. **28**, 326–329, 1969.
- [48] Nashine, H. K., Kadelburg, Z. and Radenović, S. *Common fixed point theorems for weakly isotone increasing mappings in ordered partial metric spaces*, Math. Comput. Modelling, (in press), doi:10.1016/j.mcm.2011.12.019.
- [49] Nieto, J.J. and Rodríguez-Lopez, R. *Contractive mapping theorems in partially ordered sets and applications to ordinary differential equation*, Order, **22**(3), 223–239, 2005.
- [50] Oltra, S. and Valero, O. *Banach's fixed point theorem for partial metric spaces*, Rend. Istit. Mat. Univ. Trieste **36**(1-2), 17–26, 2004.
- [51] O'Neill, S. J. *Two topologies are better than one*, (Tech. report, University of Warwick, 1995).
- [52] Paesano, D. and Vetro, P. *Suzuki's type characterizations of completeness for partial metric spaces and fixed points for partially ordered metric spaces*, Topology and its Applications **159** (3), 911–920, 2012.
- [53] Popa, V. *A general coincidence theorem for compatible multivalued mappings satisfying an implicit relation*, Demonstratio Math. **33**, 159–164, 2000.
- [54] Popa, V. *A general fixed point theorem for four weakly compatible mappings satisfying an implicit relation*, Filomat **19**, 45–51, 2005.
- [55] Ran, A.C.M. and Reurings, M.C.B. *A fixed point theorem in partially ordered sets and some applications to matrix equations*, Proc. Amer. Math. Soc. **132**, 1435–1443, 2004.
- [56] Rhoades, B.E. and Jungck, G. *Fixed points for set valued functions without continuity*, Indian J. pure and Appl. Math. **29**(3), 227–238, 1998.
- [57] Romaguera, S. and Schellekens, M. *Duality and quasi-normability for complexity spaces*, Appl. General Topology **3**, 91–112, 2002.
- [58] Romaguera, S. and Schellekens, M. *Partial metric monoids and semivaluation spaces*, Topology and Its Applications, **153** (5-6), 948–962, 2005.
- [59] Romaguera, S. and Valero, O. *A quantitative computational model for complete partial metric spaces via formal balls*, Mathematical Structures in Computer Science **19** (3), 541–563, 2009.
- [60] Romaguera, S. *A Kirk type characterization of completeness for partial metric spaces*, Fixed Point Theory and Applications **2010**, Art. ID 493298, 2010.
- [61] Romaguera, S. *Matkowski's type theorems for generalized contractions on (ordered) partial metric spaces*, Appl. General Topology **12** (2), 213–220, 2011.
- [62] Romaguera, S. *Fixed point theorems for generalized contractions on partial metric spaces*, Topology Appl. **159**, 194–199, 2012.
- [63] Samet, B. *Coupled fixed point theorems for a generalized Meir-Keeler contraction in partially ordered metric spaces*, Nonlinear Anal. **72**, 4508–4517, 2010.
- [64] Samet, B., Rajović, M., Lazović, R. and Stojiljković, R. *Common fixed point results for nonlinear contractions in ordered partial metric spaces*, Fixed Point Theory Appl. **2011**, 2011:71.
- [65] Schellekens, M. P. *A characterization of partial metrizable domains are quantifiable*, Theoretical Computer Science **305** (1–3), 409–432, 2003.
- [66] Schellekens, M. P. *The correspondence between partial metrics and semivaluations*, Theoretical Computer Science **315**(1), 135–149, 2004.
- [67] Shatanawi, W., Samet, B. and Abbas, M. *Coupled fixed point theorems for mixed monotone mappings in ordered partial metric spaces*, Math. Comput. Modelling **55** (3-4), 680–687, 2012.
- [68] Shobkolaei, N., Vaezpour, S.M. and Sedghi, S. *A common fixed point theorem on ordered partial metric spaces*, J. Basic. Appl. Sci. Res. **1** (12), 3433–3439, 2011.
- [69] Sintunavarat, W., Cho, Y. J. and Kumam, P. *Coupled coincidence point theorems for contractions without commutative condition in intuitionistic fuzzy normed spaces*, Fixed Point Theory and Applications **2011**, 2011:81.