

APPROXIMATION BY q -PHILLIPS OPERATORS

İsmet Yüksel*

Received 10:06:2010 : Accepted 27:10:2010

Abstract

In this study, we introduce a q -analogue of the Phillips operators and investigate approximation properties. We establish direct and local approximation theorems. We give a weighted approximation theorem. We estimate the rate of convergence of these operators for functions of polynomial growth on the interval $[0, \infty)$.

Keywords: Phillips operators, q -type operators, Rate of convergence, Weighted approximation, q -integral.

2010 AMS Classification: 41 A 25, 41 A 36.

Communicated by Alex Goncharov

1. Introduction

Phillips firstly introduced the q -analogue of Bernstein polynomials based on q -integer and q -binomial coefficients in [12]. Gupta and Finta obtain some direct results on certain q -Durrmeyer type operators in [6]. Recently, Aral and Gupta introduced Durrmeyer type modification of the q -Baskakov type operators in [1]. We aim to introduce a q -analogue of Phillips operators and to study approximation properties. Before this, we mention the following notations and formulas, which can be founded in [2, 8, 9] and [10]: For $n \in \mathbb{N}$, $0 < q < 1$ and $a, b \in \mathbb{R}$,

$$(1.1) \quad [n]_q = 1 + q + q^2 + \cdots + q^{n-1}, \quad n \in \mathbb{N} \setminus \{0\}; \quad [0]_q = 0,$$

$$(1.2) \quad [n]_q! = [1]_q [2]_q \cdots [n]_q, \quad n \in \mathbb{N} \setminus \{0\}; \quad [0]_q! = 1,$$

$$(1.3) \quad (a + b)_q^n = \prod_{j=1}^n (a + q^{j-1}b),$$

and

$$(1.4) \quad (1 + a)_q^\infty = \prod_{j=1}^{\infty} (1 + q^{j-1}a).$$

*Gazi University, Faculty of Science, Department of Mathematics, Teknikokullar, 06500 Beşevler, Ankara, Turkey. E-mail: iyuksel@gazi.edu.tr