

FUZZY STABILITY OF A FUNCTIONAL EQUATION RELATED TO INNER PRODUCT SPACES

Sun Young Jang* and Choonkil Park†‡

Received 01:09:2010 : Accepted 11:03:2011

Abstract

The fuzzy stability problems for the Cauchy quadratic functional equation and the Jensen quadratic functional equation in fuzzy Banach spaces have been investigated by Moslehian *et al.* Th. M. Rassias introduced the following equality

$$\sum_{i,j=1}^m \|x_i - x_j\|^2 = 2m \sum_{i=1}^m \|x_i\|^2, \quad \sum_{i=1}^m x_i = 0,$$

for a fixed integer $m \geq 3$. By the above equality, we define the following functional equation

$$(0.1) \quad \sum_{i,j=1}^m f(x_i - x_j) = 2m \sum_{i=1}^m f(x_i), \quad \sum_{i=1}^m x_i = 0.$$

In this paper, we prove the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces.

Keywords: Fuzzy Banach space, Functional equation related to inner product space, Generalized Hyers-Ulam stability.

2000 AMS Classification: 46 S 40, 46 C 05, 39 B 52, 26 E 50.

*Department of Mathematics, University of Ulsan, Ulsan 680–749, Republic of Korea.
E-mail: jsym@ulsan.ac.kr

†Department of Mathematics, Research Institute for Natural Sciences, Hanyang University, Seoul 133–791, Republic of Korea. E-mail: baak@hanyang.ac.kr

‡Corresponding Author.