

ON FUNCTION SPACES WITH WAVELET TRANSFORM IN $L_{\omega}^p(\mathbb{R}^d \times \mathbb{R}_+)$

Öznmur Kulak* and A. Turan Gürkanlı*†

Received 09:06:2010 : Accepted 19:04:2011

Abstract

Let ω_1 and ω_2 be weight functions on \mathbb{R}^d , $\mathbb{R}^d \times \mathbb{R}_+$, respectively. Throughout this paper, we define $D_{\omega_1, \omega_2}^{p, q}(\mathbb{R}^d)$ to be the vector space of $f \in L_{\omega_1}^p(\mathbb{R}^d)$ such that the wavelet transform $W_g f$ belongs to $L_{\omega_2}^q(\mathbb{R}^d \times \mathbb{R}_+)$ for $1 \leq p, q < \infty$, where $0 \neq g \in S(\mathbb{R}^d)$. We endow this space with a sum norm and show that $D_{\omega_1, \omega_2}^{p, q}(\mathbb{R}^d)$ becomes a Banach space. We discuss inclusion properties, and compact embeddings between these spaces and the dual of $D_{\omega_1, \omega_2}^{p, q}(\mathbb{R}^d)$. Later we accept that the variable s in the space $D_{\omega_1, \omega_2}^{p, q}(\mathbb{R}^d)$ is fixed. We denote this space by $(D_{\omega_1, \omega_2}^{p, q})_s(\mathbb{R}^d)$, and show that under suitable conditions $(D_{\omega_1, \omega_2}^{p, q})_s(\mathbb{R}^d)$ is an essential Banach Module over $L_{\omega_1}^1(\mathbb{R}^d)$. We obtain its approximate identities. At the end of this work we discuss the multipliers from $(D_{\omega_1, \omega_2}^{p, q})_s(\mathbb{R}^d)$ into $L_{\omega_1}^{\infty}(\mathbb{R}^d)$, and from $L_{\omega_1}^1(\mathbb{R}^d)$ into $(D_{\omega_1, \omega_2}^{p, q})_s(\mathbb{R}^d)$.

Keywords: Wavelet transform, Essential Banach module, Approximate identity, Compact embedding, Multipliers space.

2010 AMS Classification: 43 A 15, 43 A 22, 43 A 32.

Communicated by H. Turgay Kaptanoğlu

*Ondokuz Mayıs University, Faculty of Arts and Sciences, Department of Mathematics, 55139 Kurupelit, Samsun, Turkey.

E-mail: (Ö. Kulak) oznurn@omu.edu.tr (A. T. Gürkanlı) gurkanli@omu.edu.tr

†Corresponding Author.