

# A SCHUR TYPE THEOREM FOR ALMOST COSYMPLECTIC MANIFOLDS WITH KAEHLERIAN LEAVES

Nesip Aktan\*, Gülhan Ayar† and İmren Bektaş ‡ §

Received 05 : 07 : 2012 : Accepted 26 : 03 : 2013

## Abstract

In this study, we give a Schur type theorem for almost cosymplectic manifolds with Keahlerian leaves.

**Keywords:** Contact Manifold, Cosymplectic Manifold, Sectional Curvature

*2000 AMS Classification:* 53D10, 53C15, 53C25, 53C35

## 1. Introduction

Let  $M$  be a Riemannian manifold with curvature tensor  $R$ . The sectional curvature of a 2-plane  $\alpha$  in a tangent space  $T_P M$  is defined by  $K(\alpha, P) = R(X, Y, Y, X)$ , where  $\{X, Y\}$  is an orthonormal basis of  $T_P M$ . The classical theorem of F. Schur says that if  $M$  is a connected manifold of dimension  $n \geq 3$  and in any point  $P \in M$  the curvature  $K(\alpha, P)$  does not depend on  $\alpha \in T_P M$  then it does not depend on the point  $P$  too, i.e. it is a global constant. Such a manifold is called a manifold of constant sectional curvature. The Schur's theorem has been studied by many authors for different structures [11]. In 1989, Nobuhiro improves the Schur's theorem and gets a new version for locally symmetric spaces [10]. In 2001, Kassabov considers connected  $2n$ -dimensional almost Hermitian manifold  $M$  to be of pointwise constant antiholomorphic sectional curvature  $\nu(p)$ ,  $p \in M$  and proves that  $\nu$  is a global constant [6]. In 2006, Cho defines a contact strongly pseudo-convex  $CR$  space-form using the Tanaka-Webster connection in a way similar to the Sasakian space form and then he studies the geometry of such spaces. He presents a Schur type theorem for such structures [7]. The notion of an almost cosymplectic manifold was introduced by Goldberg and Yano in 1969, [19]. The simplest examples of such manifolds are those being the products (possibly local) of almost Kaehlerian manifolds and the real line  $\mathbb{R}$  or the circle  $S^1$ . Curvature properties of almost cosymplectic manifolds were studied mainly by Goldberg and Yano [12], Olszak [13], [14], Kirichenko [15] and Endo [16]. We

---

\*Duzce University, Department of Mathematics. E-Mail: (N. Aktan) nesipaktan@gmail.com

†Duzce University, Department of Mathematics. E-Mail: (G. Ayar) gulhanayar@gmail.com

‡Duzce University, Department of Mathematics. E-mail: (İ. Bektaş) bektasimren@hotmail.com

§Corresponding author