

FUZZY STABILITY OF A FUNCTIONAL EQUATION RELATED TO INNER PRODUCT SPACES

Sun Young Jang* and Choonkil Park†‡

Received 01:09:2010 : Accepted 11:03:2011

Abstract

The fuzzy stability problems for the Cauchy quadratic functional equation and the Jensen quadratic functional equation in fuzzy Banach spaces have been investigated by Moslehian *et al.* Th. M. Rassias introduced the following equality

$$\sum_{i,j=1}^m \|x_i - x_j\|^2 = 2m \sum_{i=1}^m \|x_i\|^2, \quad \sum_{i=1}^m x_i = 0,$$

for a fixed integer $m \geq 3$. By the above equality, we define the following functional equation

$$(0.1) \quad \sum_{i,j=1}^m f(x_i - x_j) = 2m \sum_{i=1}^m f(x_i), \quad \sum_{i=1}^m x_i = 0.$$

In this paper, we prove the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces.

Keywords: Fuzzy Banach space, Functional equation related to inner product space, Generalized Hyers-Ulam stability.

2000 AMS Classification: 46 S 40, 46 C 05, 39 B 52, 26 E 50.

*Department of Mathematics, University of Ulsan, Ulsan 680–749, Republic of Korea.
E-mail: jsym@ulsan.ac.kr

†Department of Mathematics, Research Institute for Natural Sciences, Hanyang University, Seoul 133–791, Republic of Korea. E-mail: baak@hanyang.ac.kr

‡Corresponding Author.

1. Introduction and preliminaries

The stability problem of functional equations originated from a question of Ulam [41] concerning the stability of group homomorphisms. Hyers [13] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [1] for additive mappings and by Th.M. Rassias [30] for linear mappings by considering an unbounded Cauchy difference. The paper of Th.M. Rassias [30] has had a lot of influence in the development of the *generalized Hyers-Ulam stability* of functional equations.

A generalization of the Th.M. Rassias theorem was obtained by Găvruta [12] by replacing the unbounded Cauchy difference by a general control function in the spirit of Th.M. Rassias' approach. During the last two decades a number of papers and research monographs have been published on various generalizations and applications of the generalized Hyers-Ulam stability to a number of functional equations and mappings (see [4, 7, 15], [21]–[27], [32]–[39]).

A square norm on an inner product space satisfies the parallelogram equality

$$\|x + y\|^2 + \|x - y\|^2 = 2\|x\|^2 + 2\|y\|^2.$$

The functional equation

$$f(x + y) + f(x - y) = 2f(x) + 2f(y)$$

is called a *quadratic functional equation*. The first author to treat the stability of the quadratic equation was F. Skof [40] by proving that if f is a mapping from a normed space X into a Banach space Y satisfying $\|f(x + y) + f(x - y) - 2f(x) - 2f(y)\| \leq \varepsilon$ for some $\varepsilon > 0$, then there is a unique quadratic mapping $g : X \rightarrow Y$ such that $\|f(x) - g(x)\| \leq \frac{\varepsilon}{2}$.

Cholewa [6] and Czerwik [8, 9] got important results on the generalized Hyers-Ulam stability problem for the quadratic functional equation.

A square norm on an inner product space satisfies

$$\sum_{i,j=1}^3 \|x_i - x_j\|^2 = 6 \sum_{i=1}^3 \|x_i\|^2$$

for all $x_1, x_2, x_3 \in \mathbb{R}$ with $x_1 + x_2 + x_3 = 0$ (see [31]).

From the above equality we can define the functional equation

$$h(x - y) + h(2x + y) + h(x + 2y) = 3h(x) + 3h(y) + 3h(x + y),$$

which can be also called a *quadratic functional equation*. In fact, $h(x) = ax^2$ in \mathbb{R} satisfies the above quadratic functional equation. In particular, every solution of the quadratic functional equation is said to be a *quadratic mapping*.

In [28], Park investigated the functional equation (0.1) and proved the generalized Hyers-Ulam stability of the functional equation (0.1) in real Banach spaces. In [29], Park and Jang proved the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces by using the fixed point method.

Katsaras [16] defined a fuzzy norm on a vector space to construct a fuzzy vector topological structure on the space. Some mathematicians have defined fuzzy norms on a vector space from various points of view [10, 18, 42]. In particular, Bag and Samanta [2], following Cheng and Mordeson [5], gave an idea of fuzzy norm in such a manner that the corresponding fuzzy metric is of Kramosil and Michalek type [17]. They established a decomposition theorem of a fuzzy norm into a family of crisp norms and investigated some properties of fuzzy normed spaces [3].

We use the definition of fuzzy normed spaces given in [2, 19, 20] to investigate a fuzzy version of the generalized Hyers-Ulam stability for the functional equation (0.1) in the fuzzy normed vector space setting.

1.1. Definition. [2, 19, 20] Let X be a real vector space. A function $N : X \times \mathbb{R} \rightarrow [0, 1]$ is called a *fuzzy norm* on X if for all $x, y \in X$ and all $s, t \in \mathbb{R}$,

- (N₁) $N(x, t) = 0$ for $t \leq 0$;
- (N₂) $x = 0$ if and only if $N(x, t) = 1$ for all $t > 0$;
- (N₃) $N(cx, t) = N(x, \frac{t}{|c|})$ if $c \neq 0$;
- (N₄) $N(x + y, s + t) \geq \min\{N(x, s), N(y, t)\}$;
- (N₅) $N(x, \cdot)$ is a non-decreasing function of \mathbb{R} and $\lim_{t \rightarrow \infty} N(x, t) = 1$;
- (N₆) for $x \neq 0$, $N(x, \cdot)$ is continuous on \mathbb{R} .

The pair (X, N) is called a *fuzzy normed vector space*.

The properties of fuzzy normed vector spaces and examples of fuzzy norms are given in [19, 20].

1.2. Definition. [2, 19, 20] Let (X, N) be a fuzzy normed vector space. A sequence $\{x_n\}$ in X is said to be *convergent*, or to *converge*, if there exists an $x \in X$ such that $\lim_{n \rightarrow \infty} N(x_n - x, t) = 1$ for all $t > 0$. In this case, x is called the *limit* of the sequence $\{x_n\}$ and we denote it by $N\text{-}\lim_{n \rightarrow \infty} x_n = x$.

1.3. Definition. [2, 19, 20] Let (X, N) be a fuzzy normed vector space. A sequence $\{x_n\}$ in X is called *Cauchy* if for each $\varepsilon > 0$ and each $t > 0$ there exists an $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ and all $p > 0$, we have $N(x_{n+p} - x_n, t) > 1 - \varepsilon$.

It is well-known that every convergent sequence in a fuzzy normed vector space is Cauchy. If each Cauchy sequence is convergent, then the fuzzy norm is said to be *complete* and the fuzzy normed vector space is called a *fuzzy Banach space*.

We say that a mapping $f : X \rightarrow Y$ between fuzzy normed vector spaces X and Y is continuous at a point $x_0 \in X$ if for each sequence $\{x_n\}$ converging to x_0 in X , the sequence $\{f(x_n)\}$ converges to $f(x_0)$. If $f : X \rightarrow Y$ is continuous at each $x \in X$, then $f : X \rightarrow Y$ is said to be *continuous* on X (see [3]).

This paper is organized as follows: In Section 2, we prove the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for the even case. In Section 3, we prove the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for the odd case.

Throughout this paper, assume that X is a vector space and that (Y, N) is a fuzzy Banach space.

2. Generalized Hyers-Ulam stability of the functional equation (0.1): the even case

In this section, we prove the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for the even case.

2.1. Lemma. [28] *Let V and W be real vector spaces. If a mapping $f : V \rightarrow W$ satisfies*

$$(2.1) \quad \sum_{i,j=1}^m f(x_i - x_j) = 2m \sum_{i=1}^m f(x_i)$$

for all $x_1, \dots, x_m \in V$ with $\sum_{i=1}^m x_i = 0$, then the mapping $f : V \rightarrow W$ is realized as the sum of an additive mapping and a quadratic mapping. \square

For a given mapping $f : X \rightarrow Y$, we define

$$Df(x_1, \dots, x_m) := \sum_{i,j=1}^m f(x_i - x_j) - 2m \sum_{i=1}^m f(x_i)$$

for all $x_1, \dots, x_m \in X$ with $\sum_{i=1}^m x_i = 0$.

2.2. Theorem. *Let $\varphi : X^m \rightarrow [0, \infty)$ be a function such that*

$$(2.2) \quad \tilde{\varphi}(x_1, \dots, x_m) := \sum_{j=1}^{\infty} 4^{-j} \varphi(2^j x_1, \dots, 2^j x_m) < \infty$$

for all $x_1, \dots, x_m \in X$. Let $f : X \rightarrow Y$ be an even mapping with $f(0) = 0$ such that

$$(2.3) \quad \lim_{t \rightarrow \infty} N(Df(x_1, \dots, x_m), t\varphi(x_1, \dots, x_m)) = 1$$

uniformly on X^m . Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^{-n} f(2^n x)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$(2.4) \quad N(Df(x_1, \dots, x_m), \delta \tilde{\varphi}(x_1, \dots, x_m)) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$(2.5) \quad N\left(f(x) - Q(x), \delta \tilde{\varphi}(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}})\right) \geq \alpha$$

for all $x \in X$.

Furthermore, the quadratic mapping $Q : X \rightarrow Y$ is a unique mapping such that

$$(2.6) \quad \lim_{t \rightarrow \infty} N(f(x) - Q(x), t \tilde{\varphi}(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}})) = 1$$

uniformly on X .

Proof. For a given $\varepsilon > 0$, by (2.3), we can find some $t_0 > 0$ such that

$$(2.7) \quad N(Df(x_1, \dots, x_m), t\varphi(x_1, \dots, x_m)) \geq 1 - \varepsilon$$

for all $t \geq t_0$. Letting $x_1 = x$, $x_2 = -x$ and $x_3 = \dots = x_m = 0$ in (2.7), we get

$$(2.8) \quad N\left(2f(2x) - 8f(x), t\varphi(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}})\right) \geq 1 - \varepsilon$$

for all $x \in X$. By induction on n , we will show that

$$(2.9) \quad N\left(f(2^n x) - 4^n f(x), t \sum_{k=1}^n 4^{n-k} \varphi\left(2^{k-1} x, -2^{k-1} x, \underbrace{0, \dots, 0}_{m-2 \text{ times}}\right)\right) \geq 1 - \varepsilon$$

for all $t \geq t_0$, all $x \in X$ and all $n \in \mathbb{N}$.

It follows from (2.8) that

$$N\left(f(2x) - 4f(x), t\varphi\left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}}\right)\right) \geq 1 - \varepsilon$$

for all $x \in X$. Thus we get (2.9) for $n = 1$.

Assume that (2.9) holds for $n \in \mathbb{N}$. Then

$$\begin{aligned} & N \left(4^{n+1} f(x) - f(2^{n+1}x), t \sum_{k=1}^{n+1} 4^{n-k+1} \varphi \left(2^{k-1}x, -2^{k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \\ & \geq \min \left\{ N \left(4^{n+1} f(x) - 4f(2^n x), t_0 \sum_{k=1}^n 4^{n-k} \varphi \left(2^{k-1}x, -2^{k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right), \right. \\ & \quad \left. N \left(4f(2^n x) - f(2^{n+1}x), t_0 \varphi \left(2^n x, -2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \right\} \\ & \geq \min\{1 - \varepsilon, 1 - \varepsilon\} = 1 - \varepsilon. \end{aligned}$$

This completes the induction argument. Letting $t = t_0$ and replacing n and x by p and $2^n x$ in (2.9), respectively, we get

$$(2.10) \quad N \left(\frac{f(2^n x)}{4^n} - \frac{f(2^{n+p}x)}{4^{n+p}}, \frac{t_0}{4^{n+p}} \sum_{k=1}^p 4^{p-k} \varphi \left(2^{n+k-1}x, -2^{n+k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \geq 1 - \varepsilon$$

for all integers $n \geq 0, p > 0$.

It follows from (2.2) and the equality

$$\begin{aligned} & \sum_{k=1}^p 4^{-n-k} \varphi \left(2^{n+k-1}x, -2^{n+k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \\ & = \sum_{k=n+1}^{n+p} 4^{-k} \varphi \left(2^{k-1}x, -2^{k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \end{aligned}$$

that for a given $\delta > 0$ there is an $n_0 \in \mathbb{N}$ such that

$$t_0 \sum_{k=n+1}^{n+p} 4^{-k} \varphi \left(2^{k-1}x, -2^{k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) < \delta$$

for all $n \geq n_0$ and $p > 0$. Now we deduce from (2.9) that

$$\begin{aligned} & N \left(4^{-n} f(2^n x) - 4^{-(n+p)} f(2^{n+p}x), \delta \right) \\ & \geq N \left(4^{-n} f(2^n x) - 4^{-(n+p)} f(2^{n+p}x), \right. \\ & \quad \left. \frac{t_0}{4^{n+p}} \sum_{k=1}^p 4^{p-k} \varphi \left(2^{n+k-1}x, -2^{n+k-1}x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \\ & \geq 1 - \varepsilon \end{aligned}$$

for each $n \geq n_0$ and all $p > 0$. Thus the sequence $\{4^{-n} f(2^n x)\}$ is Cauchy in Y . Since Y is a fuzzy Banach space, the sequence $\{4^{-n} f(2^n x)\}$ converges to some $Q(x) \in Y$. So we can define a mapping $Q : X \rightarrow Y$ by $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^{-n} f(2^n x)$, namely, for each $t > 0$ and $x \in X$, $\lim_{n \rightarrow \infty} N(4^{-n} f(2^n x) - Q(x), t) = 1$.

It is obvious that $Q : X \rightarrow Y$ is even, since $f : X \rightarrow Y$ is even.

Let $x_1, \dots, x_m \in X$. Fix $t > 0$ and $0 < \varepsilon < 1$. Since

$$\lim_{n \rightarrow \infty} 4^{-n} \varphi \left(2^n x, -2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) = 0,$$

there is an $n_1 > n_0$ such that $t_0 \varphi \left(2^n x, -2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) < \frac{4^n t}{(m^2 + m + 2)}$ for all $n \geq n_1$.

Hence for each $k \geq n_1$, we have

$$\begin{aligned} N(DQ(x_1, \dots, x_m), t) &= N \left(\sum_{i,j=1}^m Q(x_i - x_j) - 2m \sum_{i=1}^m Q(x_i), t \right) \\ &\geq \min_{1 \leq i, j \leq m} \left\{ N \left(Q(x_i - x_j) - 4^{-k} f(2^k x_i - 2^k x_j), \frac{t}{m^2 + m + 2} \right), \right. \\ &\quad N \left(2mQ(x_i) - 2m4^{-k} f(2^k x_i), \frac{t}{m^2 + m + 2} \right), \\ &\quad \left. N \left(Df(2^k x_1, \dots, 2^k x_m), \frac{2t}{(m^2 + m + 2)} \right) \right\}. \end{aligned}$$

The first $m^2 + m$ terms on the right-hand side of the above inequality tend to 1 as $k \rightarrow \infty$, and the last term is greater than

$$N \left(Df(2^k x_1, \dots, 2^k x_m), t_0 \varphi(2^k x_1, \dots, 2^k x_m) \right),$$

which is greater than or equal to $1 - \varepsilon$. Thus

$$N(DQ(x_1, \dots, x_m), t) \geq 1 - \varepsilon$$

for all $t > 0$. Since $N(DQ(x_1, \dots, x_m), t) = 1$ for all $t > 0$, by (N_2) , $DQ(x_1, \dots, x_m) = 0$ for all $x \in X$. By [28, Lemma 2.1], the mapping $Q : X \rightarrow Y$ is quadratic.

Now let for some positive δ and α , (2.4) hold. Let

$$\varphi_n(x_1, \dots, x_m) := \sum_{k=1}^n 4^{-k} \varphi(2^k x_1, \dots, 2^k x_m)$$

for all $x_1, \dots, x_m \in X$. Let $x \in X$. By the same reasoning as in the beginning of the proof, one can deduce from (2.4) that

$$(2.11) \quad N \left(4^n f(x) - f(2^n x), \delta \sum_{k=1}^n 4^{n-k} \varphi \left(2^k x, -2^k x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \geq \alpha$$

for all positive integers n . Let $t > 0$. We have

$$(2.12) \quad \begin{aligned} & N \left(f(x) - Q(x), \delta\varphi_n \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) + t \right) \\ & \geq \min \left\{ N \left(f(x) - 4^{-n} f(2^n x), \delta\varphi_n \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right), \right. \\ & \qquad \qquad \qquad \left. N(4^{-n} f(2^n x) - Q(x), t) \right\} \end{aligned}$$

Combining (2.11) and (2.12) and the fact that $\lim_{n \rightarrow \infty} N(4^{-n} f(2^n x) - Q(x), t) = 1$, we observe that

$$N \left(f(x) - Q(x), \delta\varphi_n \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) + t \right) \geq \alpha$$

for large enough $n \in \mathbb{N}$. Since the function $N(f(x) - Q(x), \cdot)$ is continuous, we see that

$$N \left(f(x) - Q(x), \delta\tilde{\varphi} \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) + t \right) \geq \alpha.$$

Letting $t \rightarrow 0$, we conclude that

$$N \left(f(x) - Q(x), \delta\tilde{\varphi} \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \geq \alpha.$$

To end the proof, it remains to prove the uniqueness assertion. Let T be another quadratic mapping satisfying (2.1) and (2.6). Fix $c > 0$. Given $\varepsilon > 0$, by (2.6) for Q and T , we can find some $t_0 > 0$ such that

$$\begin{aligned} & N \left(f(x) - Q(x), t\tilde{\varphi} \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \geq 1 - \varepsilon, \\ & N \left(f(x) - T(x), t\tilde{\varphi} \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \geq 1 - \varepsilon \end{aligned}$$

for all $x \in X$ and all $t \geq t_0$. Fix some $x \in X$ and find some integer n_0 such that

$$t_0 \sum_{k=n}^{\infty} 4^{-k} \varphi \left(2^k x, -2^k x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) < \frac{c}{2}$$

for all $n \geq n_0$. Since

$$\begin{aligned}
& \sum_{k=n}^{\infty} 4^{-k} \varphi \left(2^k x, -2^k x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \\
&= 4^{-n} \sum_{k=n}^{\infty} 4^{(n-k)} \varphi \left(2^{k-n} 2^n x, -2^{k-n} 2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \\
&= 4^{-n} \sum_{l=0}^{\infty} 4^{-l} \varphi \left(2^l 2^n x, -2^l 2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \\
&= 4^{-n} \tilde{\varphi} \left(2^n x, -2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right),
\end{aligned}$$

we have

$$\begin{aligned}
& N(Q(x) - T(x), c) \\
&\geq \min \left\{ N \left(4^{-n} f(2^n x) - Q(x), \frac{c}{2} \right), N \left(T(x) - 4^{-n} f(2^n x), \frac{c}{2} \right) \right\} \\
&= \min \left\{ N \left(f(2^n x) - Q(2^n x), 4^n \frac{c}{2} \right), N \left(T(2^n x) - f(2^n x), 4^n \frac{c}{2} \right) \right\} \\
&\geq \min \left\{ N \left(f(2^n x) - Q(2^n x), 4^n t_0 \sum_{k=n}^{\infty} 4^{-k} \varphi \left(2^k x, -2^k x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right), \right. \\
&\quad \left. N \left(T(2^n x) - f(2^n x), 4^n t_0 \sum_{k=n}^{\infty} 4^{-k} \varphi \left(2^k x, -2^k x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \right\} \\
&= \min \left\{ N \left(f(2^n x) - Q(2^n x), t_0 \tilde{\varphi} \left(2^n x, -2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right), \right. \\
&\quad \left. N \left(T(2^n x) - f(2^n x), t_0 \tilde{\varphi} \left(2^n x, -2^n x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \right\} \\
&\geq 1 - \varepsilon.
\end{aligned}$$

It follows that $N(Q(x) - T(x), c) = 1$ for all $c > 0$. Thus $Q(x) = T(x)$ for all $x \in X$. \square

2.3. Corollary. *Let $\theta \geq 0$ and let p be a real number with $p > 2$. Let $f : X \rightarrow Y$ be an even mapping such that*

$$(2.13) \quad \lim_{t \rightarrow \infty} N \left(Df(x_1, \dots, x_m), t\theta \sum_{i=1}^m \|x_i\|^p \right) = 1$$

uniformly on X^m . Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^{-n} f(2^n x)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N \left(Df(x_1, \dots, x_m), \delta\theta \sum_{i=1}^m \|x_i\|^p \right) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N \left(f(x) - Q(x), \frac{2^p}{2^p - 4} \delta \theta \|x\|^p \right) \geq \alpha$$

for all $x \in X$.

Furthermore, the quadratic mapping $Q : X \rightarrow Y$ is the unique mapping such that

$$\lim_{t \rightarrow \infty} N \left(f(x) - Q(x), \frac{2^p}{2^p - 4} t \theta \|x\|^p \right) = 1$$

uniformly on X .

Proof. Define $\varphi(x_1, \dots, x_m) := \theta \sum_{i=1}^m (\|x_i\|^p)$ and apply Theorem 2.2 to get the result, as desired. \square

Similarly, we can obtain the following. We will omit the proof.

2.4. Theorem. Let $\varphi : X^m \rightarrow [0, \infty)$ be a function such that

$$\tilde{\varphi}(x_1, \dots, x_m) := \sum_{n=0}^{\infty} 4^n \varphi \left(\frac{x_1}{2^n}, \dots, \frac{x_m}{2^n} \right) < \infty$$

for all $x_1, \dots, x_m \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.3) and $f(0) = 0$. Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^n f(\frac{x}{2^n})$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N(Df(x_1, \dots, x_m), \delta \varphi(x_1, \dots, x_m)) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N \left(f(x) - Q(x), \delta \tilde{\varphi} \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) \geq \alpha$$

for all $x \in X$.

Furthermore, the quadratic mapping $Q : X \rightarrow Y$ is the unique mapping such that

$$\lim_{t \rightarrow \infty} N \left(f(x) - Q(x), t \tilde{\varphi} \left(x, -x, \underbrace{0, \dots, 0}_{m-2 \text{ times}} \right) \right) = 1$$

uniformly on X . \square

2.5. Corollary. Let $\theta \geq 0$ and let p be a real number with $0 < p < 2$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.13). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^n f(\frac{x}{2^n})$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N \left(Df(x_1, \dots, x_m), \delta \theta \sum_{i=1}^m (\|x_i\|^p) \right) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N \left(f(x) - Q(x), \frac{2^p}{4 - 2^p} \delta \theta \|x\|^p \right) \geq \alpha$$

for all $x \in X$.

Furthermore, the quadratic mapping $Q : X \rightarrow Y$ is the unique mapping such that

$$\lim_{t \rightarrow \infty} N \left(f(x) - Q(x), \frac{2^p}{4 - 2^p} t \theta \|x\|^p \right) = 1$$

uniformly on X .

Proof. Define $\varphi(x_1, \dots, x_m) := \theta \sum_{i=1}^m (\|x_i\|^p)$ and apply Theorem 2.4 to get the result, as desired. \square

3. Generalized Hyers-Ulam stability of the functional equation (0.1): the odd case

In this section, we prove the generalized Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for the odd case.

3.1. Theorem. Let $\varphi : X^m \rightarrow [0, \infty)$ be a function such that

$$\tilde{\varphi}(x_1, \dots, x_m) := \sum_{j=1}^{\infty} 2^j \varphi\left(\frac{x_1}{2^j}, \dots, \frac{x_m}{2^j}\right) < \infty$$

for all $x_1, \dots, x_m \in X$. Let $f : X \rightarrow Y$ be an odd mapping such that

$$(3.1) \quad \lim_{t \rightarrow \infty} N(Df(x_1, \dots, x_m), t\tilde{\varphi}(x_1, \dots, x_m)) = 1$$

uniformly on X^m . Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N(Df(x_1, \dots, x_m), \delta\tilde{\varphi}(x_1, \dots, x_m)) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N\left(f(x) - A(x), \delta\tilde{\varphi}(x, x, -2x, \underbrace{0, \dots, 0}_{m-3 \text{ times}})\right) \geq \alpha$$

for all $x \in X$.

Furthermore, the additive mapping $A : X \rightarrow Y$ is a unique mapping such that

$$\lim_{t \rightarrow \infty} N(f(x) - A(x), t\tilde{\varphi}(x, x, -2x, \underbrace{0, \dots, 0}_{m-3 \text{ times}})) = 1$$

uniformly on X .

Proof. For a given $\varepsilon > 0$, by (3.1), we can find some $t_0 > 0$ such that

$$(3.2) \quad N(Df(x_1, \dots, x_m), t\varphi(x_1, \dots, x_m)) \geq 1 - \varepsilon$$

for all $t \geq t_0$. Letting $x_1 = x, x_2 = x, x_3 = -2x$ and $x_3 = \dots = x_m = 0$ in (3.2), we get

$$N\left(f(2x) - 2f(x), t\varphi(x, x, -2x, \underbrace{0, \dots, 0}_{m-3 \text{ times}})\right) \geq 1 - \varepsilon$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.2. \square

3.2. Corollary. Let $\theta \geq 0$ and let p be a real number with $p > 1$. Let $f : X \rightarrow Y$ be an odd mapping such that

$$(3.3) \quad \lim_{t \rightarrow \infty} N\left(Df(x_1, \dots, x_m), t\theta \sum_{i=1}^m \|x_i\|^p\right) = 1$$

uniformly on X^m . Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N\left(Df(x_1, \dots, x_m), \delta\theta \sum_{i=1}^m \|x_i\|^p\right) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N \left(f(x) - A(x), \frac{2^p}{2^p - 2} \delta \theta \|x\|^p \right) \geq \alpha$$

for all $x \in X$.

Furthermore, the additive mapping $A : X \rightarrow Y$ is a unique mapping such that

$$\lim_{t \rightarrow \infty} N \left(f(x) - A(x), \frac{2^p}{2^p - 2} t \theta \|x\|^p \right) = 1$$

uniformly on X .

Proof. Define $\varphi(x_1, \dots, x_m) := \theta \sum_{i=1}^m (\|x_i\|^p)$ and apply Theorem 3.1 to get the result, as desired. \square

Similarly, we can obtain the following. We will omit the proof.

3.3. Theorem. Let $\varphi : X^m \rightarrow [0, \infty)$ be a function such that

$$\tilde{\varphi}(x_1, \dots, x_m) := \sum_{n=0}^{\infty} 2^{-n} \varphi(2^n x_1, \dots, 2^n x_m) < \infty$$

for all $x_1, \dots, x_m \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (3.1). Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N(Df(x_1, \dots, x_m), \delta \varphi(x_1, \dots, x_m)) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N \left(f(x) - A(x), \delta \tilde{\varphi} \left(x, x, -2x, \underbrace{0, \dots, 0}_{m-3 \text{ times}} \right) \right) \geq \alpha$$

for all $x \in X$.

Furthermore, the additive mapping $A : X \rightarrow Y$ is a unique mapping such that

$$\lim_{t \rightarrow \infty} N \left(f(x) - A(x), t \tilde{\varphi} \left(x, x, -2x, \underbrace{0, \dots, 0}_{m-3 \text{ times}} \right) \right) = 1$$

uniformly on X .

3.4. Corollary. Let $\theta \geq 0$ and let p be a real number with $0 < p < 1$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (3.3). Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} 2^{-n} f(2^n x)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that if for some $\delta > 0, \alpha > 0$

$$N \left(Df(x_1, \dots, x_m), \delta \theta \sum_{i=1}^m (\|x_i\|^p) \right) \geq \alpha$$

for all $x_1, \dots, x_m \in X$, then

$$N \left(f(x) - A(x), \frac{2^p}{2 - 2^p} \delta \theta \|x\|^p \right) \geq \alpha$$

for all $x \in X$.

Furthermore, the additive mapping $A : X \rightarrow Y$ is a unique mapping such that

$$\lim_{t \rightarrow \infty} N \left(f(x) - A(x), \frac{2^p}{2 - 2^p} t \theta \|x\|^p \right) = 1$$

uniformly on X .

Proof. Define $\varphi(x_1, \dots, x_m) := \theta \sum_{i=1}^m (\|x_i\|^p)$ and apply Theorem 3.3 to get the result, as desired. \square

Acknowledgement

The first author and the second author were supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (NRF-2010-0013211) and (NRF-2009-0070788), respectively. This paper was written during the first author's visit to the Research Institute of Mathematics, Seoul National University.

References

- [1] Aoki, T. *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2**, 64–66, 1950.
- [2] Bag, T. and Samanta, S. K. *Finite dimensional fuzzy normed linear spaces*, J. Fuzzy Math. **11**, 687–705, 2003.
- [3] Bag, T. and Samanta, S. K. *Fuzzy bounded linear operators*, Fuzzy Sets and Systems **151**, 513–547, 2005.
- [4] Baktash, E., Cho, Y. J., Jalili, M., Saadati, R. and Vaezpour, S. M. *On the stability of cubic mappings and quadratic mappings in random normed spaces*, J. Inequal. Appl. **2008**, Art. ID 902187, 2008.
- [5] Cheng, S. C. and Mordeson, J. M. *Fuzzy linear operators and fuzzy normed linear spaces*, Bull. Calcutta Math. Soc. **86**, 429–436, 1994.
- [6] Cholewa, P. W. *Remarks on the stability of functional equations*, Aequationes Math. **27**, 76–86, 1984.
- [7] Czerwik, P. *Functional Equations and Inequalities in Several Variables* (World Scientific Publishing Company, New Jersey, Hong Kong, Singapore and London, 2002).
- [8] Czerwik, S. *On the stability of the quadratic mapping in normed spaces*, Abh. Math. Sem. Univ. Hamburg **62**, 59–64, 1992.
- [9] Czerwik, S. *The stability of the quadratic functional equation*, in: (Stability of mappings of Hyers-Ulam type, Hadronic Press, Palm Harbor, FL, 1994), 81–91.
- [10] Felbin, C. *Finite dimensional fuzzy normed linear spaces*, Fuzzy Sets and Systems **48**, 239–248, 1992.
- [11] Gajda, Z. *On stability of additive mappings*, Internat. J. Math. Math. Sci. **14**, 431–434, 1991.
- [12] Găvruta, P. *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **184**, 431–436, 1994.
- [13] Hyers, D. H. *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27**, 222–224, 1941.
- [14] Hyers, D. H., Isac, G. and Rassias, Th. M. *Stability of Functional Equations in Several Variables* (Birkhäuser, Basel, 1998).
- [15] Jung, S. *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis* (Hadronic Press Inc., Palm Harbor, Florida, 2001).
- [16] Katsaras, A. K. *Fuzzy topological vector spaces II*, Fuzzy Sets and Systems **12**, 143–154, 1984.
- [17] Kramosil, I. and Michalek, J. *Fuzzy metric and statistical metric spaces*, Kybernetika **11**, 326–334, 1975.
- [18] Krishna, S. V. and Sarma, K. K. M. *Separation of fuzzy normed linear spaces*, Fuzzy Sets and Systems **63**, 207–217, 1994.
- [19] Mirmostafae, A. K., Mirzavaziri, M. and Moslehian, M. S. *Fuzzy stability of the Jensen functional equation*, Fuzzy Sets and Systems **159**, 730–738, 2008.
- [20] Mirmostafae, A. K. and Moslehian, M. S. *Fuzzy versions of Hyers-Ulam-Rassias theorem*, Fuzzy Sets and Systems **159**, 720–729, 2008.
- [21] Park, C. *On the stability of the linear mapping in Banach modules*, J. Math. Anal. Appl. **275**, 711–720, 2002.

- [22] Park, C. *Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras*, Fixed Point Theory and Applications **2007**, Art. ID 50175, 2007.
- [23] Park, C. *Generalized Hyers-Ulam-Rassias stability of quadratic functional equations: a fixed point approach*, Fixed Point Theory and Applications **2008**, Art. ID 493751, 2008.
- [24] Park, C. and Cui, J. *Generalized stability of C^* -ternary quadratic mappings*, Abstract Appl. Anal. **2007**, Art. ID 23282, 2007.
- [25] Park, C. and Najati, A. *Homomorphisms and derivations in C^* -algebras*, Abstract Appl. Anal. **2007**, Art. ID 80630, 2007.
- [26] Park, C., Cho, Y. and Han, M. *Functional inequalities associated with Jordan-von Neumann type additive functional equations*, J. Inequal. Appl. **2007**, Art. ID 41820, 2007.
- [27] Park, C., Park, W. and Najati, A. *Functional equations related to inner product spaces*, Abstract Appl. Anal. **2009**, Art. ID 907121, 2009.
- [28] Park, C. *Fixed points, inner product spaces and functional equations* (preprint).
- [29] Park, C. and Jang, S. *Fuzzy stability of functional equations induced by inner product: a fixed point approach* (preprint).
- [30] Rassias, Th. M. *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72**, 297–300, 1978.
- [31] Rassias, Th. M. *On characterizations of inner product spaces and generalizations of the H. Bohr inequality*, in *Topics in Mathematical Analysis* (ed. Th.M. Rassias) (World Scientific Publ. Co., Singapore, 1989), 803–819.
- [32] Rassias, Th. M. *On the stability of the quadratic functional equation and its applications*, Studia Univ. Babeş-Bolyai **XLIII**, 89–124, 1998.
- [33] Rassias, Th. M. *The problem of S. M. Ulam for approximately multiplicative mappings*, J. Math. Anal. Appl. **246**, 352–378, 2000.
- [34] Rassias, Th. M. *On the stability of functional equations in Banach spaces*, J. Math. Anal. Appl. **251**, 264–284.
- [35] Rassias, Th. M. *On the stability of functional equations and a problem of Ulam*, Acta Appl. Math. **62**, 23–130, 2000.
- [36] Rassias, Th. M. *Problem 16; 2*, Report of the 27th International Symp. on Functional Equations, Aequationes Math. **39**, 292–293; 309, 1990.
- [37] Rassias, Th. M. and Šemrl, P. *On the behaviour of mappings which do not satisfy Hyers-Ulam stability*, Proc. Amer. Math. Soc. **114**, 989–993, 1992.
- [38] Rassias, Th. M. and Šemrl, P. *On the Hyers-Ulam stability of linear mappings*, J. Math. Anal. Appl. **173**, 325–338, 1993.
- [39] Rassias, Th. M. and Shibata, K. *Variational problem of some quadratic functionals in complex analysis*, J. Math. Anal. Appl. **228**, 234–253, 1998.
- [40] Skof, F. *Proprietà locali e approssimazione di operatori*, Rend. Sem. Mat. Fis. Milano **53**, 113–129, 1983.
- [41] Ulam, S. M. *A Collection of the Mathematical Problems* (Interscience Publ., New York, 1960).
- [42] Xiao, J. Z. and Zhu, X. H. *Fuzzy normed spaces of operators and its completeness*, Fuzzy Sets and Systems **133**, 389–399, 2003.