

## A RECURRENCE RELATION FOR BERNOULLI NUMBERS

Ömer Küçükşakallı\*

Received 11:06:2012 : Accepted 14:01:2013

### Abstract

Inspired by a result of Saalschütz, we prove a recurrence relation for Bernoulli numbers. This recurrence relation has an interesting connection with real cyclotomic fields.

**Keywords:** Bernoulli numbers, cyclotomic fields, Newton's identities.

*2000 AMS Classification:* 11B68, 11R18

### 1. Introduction

The Bernoulli numbers  $B_n$ , which can be defined by the Laurent series expansion

$$\frac{x}{e^x - 1} = \sum_{j=0}^{\infty} \frac{B_j}{j!} x^j,$$

have several applications in different fields of mathematics, such as number theory, combinatorics, numerical analysis.

A classical problem in elementary number theory is to find formulas for summing the  $m$ -th powers of the first  $n - 1$  integers. The following result is obtained by Bernoulli [7, Chap. 15] and it is one of the most historical formula including Bernoulli numbers

$$(1.1) \quad \sum_{j=1}^{n-1} j^m = \frac{1}{m+1} \sum_{j=0}^m \binom{m+1}{j} B_j n^{m+1-j}.$$

Another historical formula including Bernoulli numbers is the Euler-Maclaurin summation formula which was found by Euler and Maclaurin independently in 1730s and used for computations in numerical analysis [2], [9]. A breakthrough in algebraic number theory is the Kummer's criterion, proved in 1850, which relates the numerator of Bernoulli numbers to the existence of integer solutions of Fermat's equation [8].

It is a powerful method to investigate Bernoulli numbers by recurrence relations. For example, in order to prove von Staudt-Clausen theorem, concerning the denominator of the Bernoulli numbers, it is important to write  $B_j$  in terms of previous Bernoulli numbers

---

\*Mathematics Department, Middle East Technical University, 06531, Ankara, Turkey Email: [komer@metu.edu.tr](mailto:komer@metu.edu.tr)